

Cancer risk among adolescents and young adults living with HIV in South Africa: a national cohort study

Yann Ruffieux¹, Tafadzwa Dhokotera^{1,2,3,7}, Mazvita Muchengeti^{2,4}, Lina Bartels¹, Victor Olago², Julia
Bohlius^{1,3}, Elvira Singh^{2,4}, Matthias Egger^{1,5,6}, Eliane Rohner¹

Affiliations

1. Institute of Social and Preventive Medicine (ISPM), University of Bern, Switzerland
2. National Cancer Registry, National Health Laboratory Service, South Africa
3. Swiss Tropical and Public Health Institute, University of Basel, Switzerland
4. School of Public Health, University of the Witwatersrand, Johannesburg, South Africa
5. Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK
6. Centre for Infectious Disease Epidemiology and Research (CIDER), School of Public Health and Family Medicine, University of Cape Town, South Africa
7. Graduate School for Cellular and Biomedical Sciences, University of Bern, Switzerland

Correspondance to: Eliane Rohner, Institute of Social and Preventive Medicine, University of Bern, Mittelstrasse 43, 3012 Bern, Switzerland, email: eliane.rohner@ispm.unibe.ch

Abstract

Background: We studied the incidence of and risk factors for various types of cancers in adolescents and young adults living with HIV (AYALWH) in South Africa between 2004 and 2014.

Methods: We included individuals aged 15 to 24 years from the South African HIV Cancer Match study, a large cohort resulting from a linkage between HIV-related laboratory measurements from the National Health Laboratory Services and records from the National Cancer Registry. We computed incidence rates for the most common cancers. We assessed associations between these cancers and sex, age, calendar year, and CD4 cell count using Cox models and adjusted hazard ratios (aHR).

Findings: We included 782,454 AYALWH (89% female). Of those, 867 developed incident cancer including 429 who developed Kaposi sarcoma, 107 non-Hodgkin lymphoma, 48 Hodgkin lymphoma, 45 cervical cancer, and 32 leukaemia. Kaposi sarcoma was more common in the 20-24 year age group than the 15-19 year age group (aHR 1.39, 95% CI 1.03-1.86). Male sex was associated with higher rates of Kaposi sarcoma (aHR 2.06, 95% CI 1.61-2.63), non-Hodgkin lymphoma (aHR 3.17, 95% CI 2.06-4.89), Hodgkin lymphoma (aHR 4.83, 95% 2.61-8.93), and leukaemia (aHR 5.90, 95% CI 2.87-12.1). Lower CD4 cell counts at baseline were associated with higher rates of Kaposi sarcoma, cervical cancer, non-Hodgkin and Hodgkin lymphoma.

Interpretation: Infection-related cancers are the most common cancer types among AYALWH in South Africa. The burden of these cancers may be reduced through HPV vaccination, targeted HIV testing, early initiation of antiretroviral therapy, and improvement of treatment adherence.

Introduction

Adolescents and young adults in South Africa are disproportionately affected by HIV. In 2019, one in three new HIV infections in South Africa occurred among people aged 15-24 years.¹ Girls and young women are particularly vulnerable, accounting for nearly 80% of new HIV infections in this age group.¹ Compared to other age groups, adolescents living with HIV are less likely to initiate antiretroviral therapy (ART) and more likely to drop out of care and have poorly controlled HIV infection with higher rates of virologic failure.^{2,3}

HIV-induced immunodeficiency combined with a higher susceptibility to oncogenic viruses increases the risk of cancer among people living with HIV.⁴ Cancer is rare among adolescents and young adults, but an elevated cancer risk has been documented among those living with HIV, with the AIDS-defining cancers, Kaposi sarcoma and non-Hodgkin lymphoma, being the most frequently diagnosed cancers.^{5,6} However, analyses of cancer incidence and associated risk factors in this age group are often limited by small numbers of cancer cases.⁵ Furthermore, data from sub-Saharan Africa, where most young people living with HIV reside, are scarce.

The South African HIV Cancer Match (SAM) study is a nationwide cohort of people living with HIV in South Africa. We linked HIV-related laboratory records from the National Health Laboratory Services (NHLS) to cancer records from the National Cancer Registry (NCR) to create this cohort.⁷ The SAM study provided us with a unique data source to assess the incidence and spectrum of cancers among young people living with HIV in South Africa.

Methods

The SAM study

As described in detail elsewhere,⁷ the SAM study was constructed using data from the NHLS, the largest diagnostic pathology service in South Africa, and the pathology-based NCR for the years 2004 to 2014. The NHLS maintains laboratories in all nine South African provinces and stores data at a corporate data warehouse. The NHLS is estimated to provide diagnostic pathology services to approximately 80% of the South African population. We used privacy-preserving probabilistic record linkage methods to, first, identify HIV-related laboratory records which most likely belonged to the same individual and, second, link these individuals to cancer diagnoses from the NCR.⁸ HIV-related laboratory measurements included HIV tests, CD4 cell counts and percentages, and HIV RNA viral loads. Ethical approval was obtained from the Human Research Ethics Committee of the University of the Witwatersrand, Johannesburg (M190594), and the Cantonal Ethics committee in Bern (2016–00589).

Inclusion criteria and definitions

We included individuals aged between 15 and 24 years at baseline who had at least two HIV-related laboratory measurements on separate days. We defined an individual's baseline date as the date of their first positive HIV test or HIV-related measurement (CD4 cell count or RNA viral load). We excluded individuals with missing information on sex or age. Adolescents and young adults diagnosed with cancer before their baseline date were excluded from the analysis of that specific cancer.

We classified cancers based on the International Classification of Diseases for Oncology, 3rd Edition (ICD-0-3), recoded for adolescent and young adults (AYA).⁹ We categorised cancers into the following groups¹⁰: leukaemia (AYA recode 01-04), non-Hodgkin lymphoma (05), Hodgkin lymphoma (06), central nervous system (CNS) and other intracranial and intraspinal neoplasms (07-16), osseous & chondromatous neoplasms (17-20), Kaposi sarcoma (24), soft tissue sarcomas other than Kaposi sarcoma (21-23,25), germ cell and trophoblastic neoplasms (26-28), melanoma and skin carcinomas (29, 30), carcinomas (31-48), and miscellaneous specified and unspecified neoplasms and unclassified cancers (49-56, 99). We divided the non-Hodgkin lymphomas into Burkitt lymphomas (ICD-0-3 morphology code 9687) and other non-Hodgkin lymphomas. We distinguished between the following carcinomas: thyroid carcinoma, other carcinomas of the head and neck, carcinoma of the genitourinary tract other than cervix, carcinoma of the cervix (ICD-0-3 topography code C53), carcinoma of the gastrointestinal tract, and carcinoma of other and ill-defined sites.

Statistical analysis

We produced descriptive statistics of AYA living with HIV (AYALWH), overall and stratified by baseline age (15 to 19 years versus 20 to 24 years). Calendar year and CD4 cell count (cells/ μ L) were assessed at baseline. CD4 cell count was labelled as missing if no CD4 cell count measurement was available for a given individual within the first two weeks of the baseline date. We used two-sided corrected tests to compare the proportion of individuals who developed a given type of cancer between ages 15 to 19 years to the proportion of individuals who developed that type of cancer from ages 20 to 24 years, among AYALWH who developed cancer of any type.

Individuals contributed person-years starting from their baseline date to six months after their last HIV-related laboratory measurement, their 25th birthday, database closure (1st January 2015), or the first diagnosis date of the cancer(s) under consideration, whichever came first. We performed survival analyses for the following cancer groups: cancer of any type, Kaposi sarcoma, cervical cancer, non-Hodgkin lymphoma, Hodgkin lymphoma, and leukaemia. We computed crude incidence rates for these cancers by dividing the number of cases by the number of person-years at risk. The rates were

computed for ages 15 to 19 years, 20 to 24 years, and overall, treating age as a time-updated variable. We produced 95% confidence intervals (CI) for these rates assuming a Poisson distribution.

We calculated adjusted hazard ratios (aHR) using multivariable Cox proportional hazards models. These models were adjusted for age (time-updated categorical variable: 15-19 years, 20-24 years), sex, baseline calendar period (categorical variable: 2004-2007, 2008-2011, 2012-2014), and baseline CD4 cell count (categorical variable: <200 cells/ μ L, 200-499 cells/ μ L, \geq 500 cells/ μ L, missing). We assessed the proportional hazards assumption using Schoenfeld residuals. The proportional hazards assumption for the CD4 cell count was violated when analysing the 'any cancer' category. We, therefore, stratified the corresponding Cox model for that variable and estimated cumulative hazard functions and 95% confidence intervals for each CD4 cell count group using the Nelson-Aalen method. For cervical cancer, the proportional hazards assumption was violated for age group, and we stratified the corresponding Cox model with respect to that variable. Due to the limited number of cases, we only present non-adjusted hazard ratios for leukaemia. In a separate analysis, we modelled the relationship between baseline CD4 count (treated as a continuous variable) and the log-hazard using a penalised spline transformation. We excluded patients with missing CD4 cell count at baseline from this analysis. We produced aHR curves with 95% CIs over a grid of CD4 values, with a reference CD4 cell count of 200 cells/ μ L. All analyses were done in R 3.6.2 (R Foundation for Statistical Computing, Vienna, Austria).

Role of funding source

The funders of this study had no role in the design, data collection, data analysis, data interpretation, or writing of the report.

Results

Study population and cancer cases

We included 782,454 AYALWH over a study period of 11 years: 176,019 were diagnosed with HIV between the ages of 15 to 19 years and 606,435 between the ages of 20 to 24 years. The majority (89.2%; n = 698,066) of the included AYALWH were women ([Table 1](#)). The median CD4 cell count at baseline was 386 cells/ μ L (interquartile range [IQR] 253-548) in the 15-19 age group and 349 cells/ μ L (IQR 216-511) in the 20-24 age group.

Over 1,428,114 person-years, 867 AYALWH developed incident cancer between the age of 15 to 24 years ([Table 2](#)). The median follow-up time was 1.42 years (IQR 0.70-2.61). The most common cancer types were Kaposi sarcoma (429 cases) and non-Hodgkin lymphoma (107 cases), followed by Hodgkin lymphoma (48 cases), cervical cancer (45 cases), and leukaemia (32 cases). Among AYALWH who

developed cancer, the proportion of patients who developed a non-cervical carcinoma was higher in the 20-24 age group than in the 15-19 age group (13% vs. 5%, $p=0.03$).

The characteristics of AYALWH who developed cancer and those who remained free of cancer are shown in [Table 3](#). Among AYALWH who remained free of cancer, only 11% were male, but this percentage increased to 33% among those who developed Hodgkin lymphoma and 38% among those who developed leukaemia. AYALWH who developed Kaposi sarcoma, cervical cancer, non-Hodgkin lymphoma or Hodgkin lymphoma had median baseline CD4 cell counts below 300 cells/ μL . Median baseline CD4 cell counts were higher in AYALWH who developed leukaemia (385 cells/ μL , IQR 208-806) than those who remained free of cancer (357 cells/ μL , IQR 224-520).

Cancer incidence rates and risk factors

The overall cancer incidence rate among AYALWH aged 15-24 years was 60.7/100,000 person-years ([Table 4](#)). Cancer-specific incidence rates were highest for Kaposi sarcoma (30.0/100,000 person-years, 95% CI 27.2-33.0) followed by non-Hodgkin lymphoma (7.5/100,000 person-years, 95% CI 6.1-9.0). [Table 5](#) shows hazard ratios for the risk of developing various types of cancers. The risk of developing any cancer was higher in 20-24-year-olds than 15-19-year-olds (aHR 1.32, 95% CI 1.08-1.61). Similarly, the incidence rate for Kaposi sarcoma was higher in 20-24-year-olds than 15-19-year-olds (aHR 1.39, 95% CI 1.03-1.86). For cervical cancer, the incidence rate also tended to be higher in the older age group (3.8/100,000 person-years, 95% CI 2.7-5.1) than the younger age group (1.5/100,000 person-years, 95% CI 0.3-4.4, [Table 4](#)). However, because the proportional hazards assumption was violated we did not compute a hazard ratio for the association of age group with cervical cancer risk. There was no evidence of an association between age group and the incidence rates of the other types of cancers among AYALWH. The risk of developing cancer decreased in more recent baseline calendar years. Male sex was associated with higher incidence rates of any cancer (aHR 2.16, 95% CI 1.82-2.56), Kaposi sarcoma (aHR 2.06, 95% CI 1.61-2.63), non-Hodgkin lymphoma (aHR 3.17, 95% CI 2.06-4.89), Hodgkin lymphoma (aHR 4.83, 95% CI 2.61-8.93), and leukaemia (unadjusted hazard ratio 5.90, 95% CI 2.87-12.1).

The risk of developing cervical cancer decreased with increasing baseline CD4 cell count (≥ 500 versus < 200 cells/ μL , aHR 0.21, 95% CI 0.07-0.57). A similar pattern of decreasing incidence rates with higher baseline CD4 cell counts was observed for non-Hodgkin lymphoma (≥ 500 versus < 200 cells/ μL , aHR 0.30, 95% CI 0.16-0.55). Individuals with a missing CD4 cell count had a lower risk of developing Kaposi sarcoma compared to individuals with a CD4 cell count below 200 cells/ μL , but a higher risk than individuals with a CD4 cell count above 500 cells/ μL . [Supplementary Figure 1](#) shows the cumulative hazards for developing any cancer, stratified by CD4 cell count category. The cumulative hazard was

consistently highest in the group with 0-199 CD4 cells/ μ L and lowest in those with \geq 500 CD4 cells/ μ L. In [Figure 1](#), we show aHRs for the incidence of various types of cancers, comparing baseline CD4 cell count (treated as a continuous variable) to a reference value of 200 cells/ μ . We excluded 74,937 AYALWH from this analysis due to missing CD4 cell counts at baseline. A lower baseline CD4 cell count was associated with higher rates of any cancer, Kaposi Sarcoma, cervical cancer, non-Hodgkin lymphoma, and Hodgkin lymphoma, with the association being strongest for Kaposi sarcoma. There was no evidence for an association between lower CD4 cell counts and a higher risk of leukaemia.

Discussion

Kaposi sarcoma and non-Hodgkin lymphoma were the most frequent incident cancers among AYALWH in South Africa. The overall risk of developing cancer increased slightly with age and was higher in male than female AYALWH. Lower CD4 cell counts at baseline were associated with an elevated risk of Kaposi sarcoma, cervical cancer, non-Hodgkin lymphoma, and Hodgkin lymphoma after adjusting for sex and baseline calendar period.

Our analysis is one of the first large-scale studies to assess cancer risk among AYALWH and included data of more than 700,000 AYALWH and 867 incident cancer cases across South Africa. Our large sample size enabled us to separately assess incidence rates and risk factors for the most common cancer types. We censored AYALWH at their 25th birthday and treated age as a time-updated variable to describe the burden of cancers diagnosed between age 15 to 24 years. We did not include cancer diagnoses that AYALWH developed later in life. Several limitations must be noted. Even though our study of cancer risk among AYALWH may be one of the largest to date, case numbers for many specific cancer types were still small. The number of cancer cases among AYALWH is likely to be underestimated in our study, as cancer diagnoses were from the pathology-based NCR, and clinically diagnosed cases may have been missed. About two-thirds of the included AYALWH did not have an HIV viral load measurement recorded during the study period. Thus, we did not assess the association between HIV viremia and cancer risk in this population. Furthermore, as the SAM study is based on laboratory records from the NHLS, we did not have access to other relevant patient-level information such as socio-economic and behavioural data, mode of HIV infection, ART prescriptions, and human papillomavirus (HPV) vaccination status. We did not have information on the death or migration of individuals within the SAM study. Hence, we censored time-at-risk for all individuals six months after their last HIV-related laboratory measurement. We defined incident cancer cases as cancers diagnosed after the first HIV-related laboratory measurement, with no lag time. However, estimates of cancer incidence rates and hazard ratios are sensitive to the definition of incident cancers and person-time-

at-risk. Therefore, our estimates may not necessarily be comparable with estimates from other studies using different definitions.

In 2020, the most common cancer types among AYA worldwide were leukaemia, thyroid cancer and testicular cancer followed by non-Hodgkin lymphoma, CNS tumours, Hodgkin lymphoma, and breast cancer.¹¹ However, in South Africa, where close to 1 in 10 AYA are living with HIV,¹² the most frequently diagnosed cancers among AYA were the AIDS-defining cancers, Kaposi sarcoma, cervical cancer, and non-Hodgkin lymphoma.¹¹ Kaposi sarcoma was also by far the most common cancer in our study of AYALWH. This is likely due to the low proportion of ART exposure among AYALWH in South Africa,^{3,13} as ART reduces the risk of developing Kaposi sarcoma substantially.¹⁴ Furthermore, the prevalence of human herpesvirus 8 (HHV-8), the underlying infectious cause of Kaposi sarcoma, shows a distinct geographic pattern.¹⁵ In sub-Saharan Africa, where HHV-8 prevalence is high, the virus is typically transmitted horizontally during childhood.¹⁵ In Europe and North America, however, HHV-8 seems to be predominantly sexually transmitted and is particularly common among men who have sex with men.¹⁵ In line with the geographic variation of HHV-8 prevalence, crude Kaposi sarcoma incidence rates have been reported to be approximately three times higher among AYALWH (age 16-25 years) who initiated ART in South Africa compared with those who initiated ART in Europe or North America.¹⁶

Hematologic malignancies are among the most frequent cancers in AYA globally.¹¹ The risk of developing non-Hodgkin and Hodgkin lymphoma is elevated among people living with HIV, whereas leukaemia is not linked to HIV infection.¹⁷ However, studies on hematologic malignancies among AYALWH are scarce. A British study of 290 AYA aged 10-24 years who had been perinatally infected with HIV found that 6 out of 8 malignancies diagnosed in this group were lymphomas (3 Hodgkin lymphomas, 1 Burkitt lymphoma, 2 B-cell lymphomas).¹⁸ In adults, cumulative HIV viremia is a strong, independent risk factor for the development of non-Hodgkin lymphoma, particularly Burkitt lymphoma.^{19,20} For AYALWH, this is anecdotally supported by a report on two vertically infected 15-year-olds in Italy who were diagnosed with Burkitt lymphoma after prolonged exposure to HIV viremia.²¹ Compared to other age groups, AYALWH have higher rates of non-adherence to antiretroviral treatment and, thus, of virologic failure.² This may put AYALWH at particularly high risk of developing HIV-related hematologic malignancies.

Cervical cancer is one of the most preventable malignancies but still appears among the five most common cancer types in our analysis. In the general population, cervical cancer incidence rates among AYA are very low, and infections with HPV in this age group are often transient.²² Therefore, cervical cancer screening is generally not recommended for HIV-negative AYA below the age of 25 years.²³ However, HIV infection increases the risk of developing cervical cancer by a factor of six,²⁴ and, thus,

sexually active AYALWH should be screened for cervical cancer irrespective of age.²³ We found cervical cancer incidence in AYALWH to increase with lower CD4 cell counts. This is in line with what has previously been reported for adult women living with HIV.^{25,26} Timely initiation of ART and maintained adherence is likely to reduce the risk of developing cervical cancer.²⁷ Another pillar in the prevention of cervical cancer is vaccination against high-risk HPV genotypes. In South Africa, a national HPV vaccination program targeting school girls aged 9-13 years, regardless of HIV status, was launched in 2014.²⁸ Given the closing date of our analysis, most AYALWH included in the present study would not have been vaccinated against HPV yet.

Among adolescents aged 15-19 years living with HIV in the United States, 49% of cancer cases were attributable to infections.²⁹ Many of the most common cancer types among AYALWH included in our study were also infection-related: together, Kaposi sarcoma (HHV-8), non-Hodgkin and Hodgkin lymphomas (Epstein-Barr virus), and cervical cancer (HPV) made up 70% of the cancer cases. These cancer types showed a clear association with lower CD4 cell counts at baseline. HPV vaccination, cervical cancer screening, and treatment of pre-cancerous cervical lesions are vital steps towards decreasing cervical cancer incidence. Additionally, the burden of infection-related cancers among AYALWH may be reduced by preventing HIV-induced immunodeficiency through improved HIV testing and early initiation of ART. Mental health and HIV-related stigma are key issues in ALYALWH that may negatively impact treatment adherence and retention in care.³⁰

In conclusion, infection-related cancers are the most common types of cancer occurring in AYALWH in South Africa. Innovative strategies to encourage early ART initiation and enhance treatment adherence in AYALWH are needed to achieve sustained viral suppression and reduce the risk of developing cancer and other co-morbidities in this vulnerable population.

Acknowledgments and funding source

This work was supported by the National Institute of Allergy and Infectious Diseases of the National Institutes of Health (NIH) under Award Number U01AI069924. The work was also supported by the NIH administrative supplement to Existing NIH Grants and Cooperative Agreements (Parent Admin Supp) (U01AI069924-09, to ME, JB), PEPFAR supplement (to ME), the Swiss National Science Foundation (SNSF, 320030_169967, to JB, ME), and the U.S. CRDF Global (HIV_DAA3-16-62705-1, to MM). ME was supported by special project funding (17481189498) from the SNSF.

References

- 1 AIDSinfo | UNAIDS. Available from: <https://aidsinfo.unaids.org/> (accessed March 5, 2021).
- 2 Slogrove AL, Mahy M, Armstrong A, Davies MA. Living and dying to be counted: What we know about the epidemiology of the global adolescent HIV epidemic. *J Int AIDS Soc* 2017; **20**: 21520.
- 3 Maskew M, Bor J, Macleod W, Carmona S, Sherman GG, Fox MP. Adolescent HIV treatment in South Africa ' s national HIV programme : a retrospective cohort study. *Lancet HIV* 2019; **6**: e760–8.
- 4 da Silva Neto MM, Brites C, Borges ÁH. Cancer during HIV infection. *Apmis* 2020; **128**: 121–8.
- 5 Bohlius J, Foster C, Naidu G, Sengayi M, Turkova A. Cancer in adolescents and young adults living with HIV. *Curr Opin HIV AIDS* 2018; **13**: 196–203.
- 6 Dhokotera T, Bohlius J, Egger M, *et al.* Cancer in HIV-positive and HIV-negative adolescents and young adults in South Africa: a cross-sectional study. [Preprint] medRxiv. August 21, 2020. doi:10.1101/2020.08.18.20176289.
- 7 Muchengeti M, Bartels L, Olago V, *et al.* Cohort Profile: The South African HIV Cancer Match Study (SAM) [Preprint] OSF Preprints. August 11, 2020. doi:10.31219/osf.io/w52sb.
- 8 Schmidlin K, Clough-Gorr KM, Spoerri A, Group for SNC study. Privacy Preserving Probabilistic Record Linkage (P3RL): a novel method for linking existing health-related data and maintaining participant confidentiality. *BMC Med Res Methodol* 2015; **15**: 46.
- 9 Barr RD, Holowaty EJ, Birch JM. Classification schemes for tumors diagnosed in adolescents and young adults. *Cancer* 2006; **106**: 1425–30.
- 10 AYA Site Recode/WHO 2008 Definition. Available from: <https://seer.cancer.gov/ayarecode/aya-who2008.html> (accessed April 8, 2021).
- 11 Ferlay J, Ervik M, Lam F, *et al.* Global Cancer Observatory: Cancer Today. Lyon, France. International Agency for Research on Cancer. 2020. Available from: <https://gco.iarc.fr/today> (accessed March 6, 2021).
- 12 Zanoni BC, Archary M, Buchan S, Katz IT, Haberer JE. Systematic review and meta-analysis of the adolescent HIV continuum of care in South Africa: The Cresting Wave. *BMJ Glob Heal* 2016; **1**: 1.

- 13 Shisana O, Rehle T, Simbayi L, *et al.* South African National HIV Prevalence, Incidence and Behaviour Survey, 2012. Cape Town: HSRC Press, 2014. Available from: <http://www.hsrc.ac.za/en/research-data/view/6871>.
- 14 Semeere AS, Busakhala N, Martin JN. Impact of antiretroviral therapy on the incidence of Kaposi's sarcoma in resource-rich and resource-limited settings. *Curr Opin Oncol* 2012; **24**: 522–30.
- 15 Rohner E, Wyss N, Heg Z, *et al.* HIV and human herpesvirus 8 co-infection across the globe: Systematic review and meta-analysis. *Int J Cancer* 2016; **138**: 45–54.
- 16 Rohner E, Bütikofer L, Schmidlin K, *et al.* Comparison of Kaposi Sarcoma risk in human immunodeficiency virus-positive adults across 5 continents: A multiregional multicohort study. *Clin Infect Dis* 2017; **65**: 1316–26.
- 17 Kimani SM, Painschab MS, Horner MJ, *et al.* Epidemiology of haematological malignancies in people living with HIV. *Lancet HIV* 2020; **7**: e641–51.
- 18 Chhabra S, Fidler S, Ayers S, Bower M, Lyall H, Foster C. Malignancy and all-cause mortality; incidence in adolescents and young adults living with perinatally acquired HIV. *J Virus Erad* 2020; **6**: 30–3.
- 19 Zoufaly A, Stellbrink HJ, an der Heiden M, *et al.* Cumulative HIV viremia during highly active antiretroviral therapy is a strong predictor of AIDS-related lymphoma. *J Infect Dis* 2009; **200**: 79–87.
- 20 Hernández-Ramírez RU, Qin L, Lin H, *et al.* Association of immunosuppression and HIV viraemia with non-Hodgkin lymphoma risk overall and by subtype in people living with HIV in Canada and the USA: a multicentre cohort study. *Lancet HIV* 2019; **6**: e240–9.
- 21 Zangari P, Santilli V, Cotugno N, *et al.* Raising Awareness of Non-Hodgkin Lymphoma in HIV-infected Adolescents. *J Pediatr Hematol Oncol* 2013; **35**: e134–7.
- 22 Ho GYF, Bierman R, Beardsley L, Chang CJ, Burk RD. Natural History of Cervicovaginal Papillomavirus Infection in Young Women. *N Engl J Med* 1998; **338**: 423–8.
- 23 Botha M, Dreyer G. Guidelines for cervical cancer screening in South Africa. *South Afr J Gynaecol Oncol* 2017; **9**: 8–12.
- 24 Stelzle D, Tanaka LF, Lee KK, *et al.* Estimates of the global burden of cervical cancer associated with HIV. *Lancet Glob Heal* 2021; **9**: e161–9.

- 25 Ruffieux Y, Muchengeti M, Egger M, *et al.* Immunodeficiency and Cancer in 3.5 Million People Living with Human Immunodeficiency Virus: the South African HIV Cancer Match Study. *Clin Infect Dis* 2021; published online Feb 2. doi:10.1093/cid/ciab087.
- 26 Abraham AG, Strickler HD, D'Souza G. Invasive cervical cancer risk among HIV-infected women is a function of CD4 count and screening. *J Acquir Immune Defic Syndr* 2013; **63**: e163.
- 27 Kelly H, Weiss HA, Benavente Y, *et al.* Association of antiretroviral therapy with high-risk human papillomavirus, cervical intraepithelial neoplasia, and invasive cervical cancer in women living with HIV: a systematic review and meta-analysis. *Lancet HIV* 2018; **5**: e45–58.
- 28 Dlamini SK, Madhi SA, Muloiwa R, *et al.* Guidelines for the vaccination of HIV-infected adolescents and adults in South Africa. *South Afr J HIV Med* 2018; **19**: a839.
- 29 De Martel C, Shiels MS, Franceschi S, *et al.* Cancers attributable to infections among adults with HIV in the United States. *AIDS* 2015; **29**: 2173–81.
- 30 Vreeman RC, McCoy BM, Lee S. Mental health challenges among adolescents living with HIV. *J Int AIDS Soc* 2017; **20**: 100–9.

Table 1: Baseline characteristics of adolescents and young adults living with HIV and included in the analysis, stratified by age.

	Age at baseline		
	15-19 years	20-24 years	Overall
	N = 176,019	N = 606,435	N = 782,454
Sex			
Female	160,937 (91.4)	537,129 (88.6)	698,066 (89.2)
Male	15,082 (8.6)	69,306 (11.4)	84,388 (10.8)
Calendar period			
2004 – 2007	31,916 (18.1)	136,298 (22.5)	168,214 (21.5)
2008 – 2011	90,375 (51.3)	307,037 (50.6)	397,412 (50.8)
2012 – 2014	53,728 (30.5)	163,100 (26.9)	216,828 (27.7)
Median baseline CD4 cell count (IQR) [cells/μL]	386 (253-548)	349 (216-511)	357 (224-520)
Baseline CD4 cell count [cells/μL]			
<200	26,392 (15.0)	121,553 (20.0)	147,945 (18.9)
200-499	82,462 (46.8)	282,455 (46.6)	364,917 (46.6)
\geq 500	49,690 (28.2)	144,968 (23.9)	194,658 (24.9)
Missing	17,475 (9.9)	57,459 (9.5)	74,934 (9.6)

Results are presented as counts and percentages if not otherwise stated.

Table 2: Incident cancer cases in adolescents and young adults living with HIV, stratified by age at cancer diagnosis. The percentages represent the proportion of individuals who developed the specific type of cancer among those who developed any cancer, within each age group.

	Age at cancer diagnosis		
	15-19 years n (%)	20-24 years n (%)	Overall n (%)
Cancer type			
Any cancer	112 (100%)	755 (100%)	867 (100%)
Kaposi sarcoma (KS)	52 (46%)	377 (50%)	429 (50%)
Non-Hodgkin lymphoma (NHL)	18 (16%)	89 (12%)	107 (12%)
Burkitt lymphoma	2	25	27
Other NHL	16	64	80
Cervical cancer	3 (3%)	42 (6%)	45 (3%)
Leukaemia	5 (5%)	27 (4%)	32 (4%)
Hodgkin lymphoma	10 (9%)	38 (5%)	48 (6%)
CNS and other intracranial and intraspinal neoplasms	0 (0%)	2 (<1%)	2(<1%)
Osseous and chondromatous neoplasms	4 (4%)	10 (1%)	14 (2%)
Soft tissue sarcomas other than KS	6 (5%)	17 (2%)	23 (3%)
Germ cell and trophoblastic neoplasms	5 (5%)	14 (2%)	19 (2%)
Melanoma and skin carcinoma	2 (2%)	28 (4%)	30 (4%)
Melanoma	1	4	5
Skin carcinoma	1	24	25
Carcinoma excluding cervical cancer	6 (5%)	98 (13%)	104 (12%)
Thyroid cancer	0	2	2
Other head and neck cancer	0	9	9
Breast cancer	1	12	13
Genitourinary cancer other than cervix	3	25	28
Gastrointestinal carcinoma	1	22	23
Any other carcinoma	1	28	29
Miscellaneous specified and unspecified neoplasms	3 (3%)	20 (3%)	23 (3%)

Table 3: Baseline characteristics of adolescents and young adults living with HIV who developed cancer or remained free of cancer.

	Any cancer	Kaposi sarcoma	Cervical cancer	Non-Hodgkin lymphoma	Hodgkin lymphoma	Leukaemia	Free of cancer
	N=867	N=429	N=45	N=107	N=48	N=32	N=781,587
Age							
15-19 years	231 (26.6)	108 (25.2)	13 (28.9)	25 (23.4)	15 (31.2)	9 (28.1)	175,788 (22.5)
20-24 years	636 (73.4)	321 (74.8)	32 (71.1)	82 (76.6)	33 (68.8)	23 (71.9)	605,799 (77.5)
Sex							
Female	700 (80.7)	347 (80.9)	45 (100.0)	78 (72.9)	32 (66.7)	20 (62.5)	697,366 (89.2)
Male	167 (19.3)	82 (19.1)	0	29 (27.1)	16 (33.3)	12 (37.5)	167 (19.3)
Calendar period							
2004 - 2007	282 (32.5)	152 (35.4)	13 (28.9)	32 (29.9)	18 (37.5)	11 (34.4)	167,932 (21.5)
2008 - 2011	423 (48.8)	206 (48.0)	27 (60.0)	49 (45.8)	20 (41.7)	17 (53.1)	396,989 (50.8)
2012 - 2014	162 (18.7)	71 (16.6)	5 (11.1)	26 (24.3)	10 (20.8)	4 (12.5)	216,666 (27.7)
Median CD4 cell count [IQR]	248 [121, 399]	207 [84, 350]	261 [168, 361]	240 [139, 345]	267 [147, 447]	385 [208, 806]	357 [224, 520]
CD4 cell count							
<200	293 (33.8)	179 (41.7)	15 (33.3)	37 (34.6)	15 (31.2)	6 (18.8)	147,652 (18.9)
200-499	338 (39.0)	158 (36.8)	19 (42.2)	42 (39.3)	17 (35.4)	11 (34.4)	364,579 (46.6)
≥500	127 (14.6)	42 (9.8)	5 (11.1)	14 (13.1)	10 (20.8)	11 (34.4)	194,531 (24.9)
Missing	109 (12.6)	50 (11.7)	6 (13.3)	14 (13.1)	6 (12.5)	4 (12.5)	74,825 (9.6)

Results are presented as counts and percentages if not otherwise stated.

Table 4: Incidence rates for various types of cancer, stratified by current age group (time-updated).

	Incidence rate per 100,000 person-years (95% CI)		
	Age 15-19 years	Age 20-24 years	Overall
Any cancer	50.9 (41.9-61.3)	62.5 (58.1-67.1)	60.7 (56.7-64.9)
Kaposi sarcoma	23.6 (17.6-31)	31.2 (28.1-34.5)	30 (27.2-33)
Cervical cancer	1.5 (0.3-4.4)	3.8 (2.7-5.1)	3.4 (2.5-4.6)
Non-Hodgkin lymphoma	8.2 (4.8-12.9)	7.4 (5.9-9.1)	7.5 (6.1-9)
Hodgkin lymphoma	4.5 (2.2-8.4)	3.1 (2.2-4.3)	3.4 (2.5-4.5)
Leukaemia	2.3 (0.7-5.3)	2.2 (1.5-3.2)	2.2 (1.5-3.2)

CI, confidence interval.

Table 5: Adjusted hazard ratios for developing different cancer types among adolescents and young adults living with HIV.

	Hazard ratios* (95% confidence intervals)					
	Any cancer	Kaposi sarcoma	Cervical cancer	Non-Hodgkin lymphoma	Hodgkin lymphoma	Leukaemia
Age [years]						
15-19	1	1	†	1	1	1
20-24	1.32 (1.08-1.61)	1.39 (1.03-1.86)		1.05 (0.63-1.76)	0.72 (0.35-1.47)	1.19 (0.46-3.12)
Sex						
Female	1	1	-	1	1	1
Male	2.16 (1.82-2.56)	2.06 (1.61-2.63)	-	3.17 (2.06-4.89)	4.83 (2.61-8.93)	5.90 (2.87-12.1)
Calendar period						
2004 – 2007	1	1	1	1	1	1
2008 – 2011	0.71 (0.60-0.83)	0.61 (0.49-0.76)	1.25 (0.61-2.57)	0.69 (0.43-1.09)	0.56 (0.28-1.11)	0.66 (0.31-1.41)
2012 – 2014	0.69 (0.57-0.85)	0.54 (0.41-0.73)	0.72 (0.24-2.15)	0.83 (0.48-1.42)	0.61 (0.26-1.40)	0.38 (0.12-1.20)
CD4 cell count [cells/μL]						
<200		1	1	1	1	1
200-499	†	0.34 (0.28-0.42)	0.44 (0.22-0.86)	0.46 (0.29-0.72)	0.47 (0.23-0.95)	0.67 (0.25-1.81)
≥500		0.17 (0.12-0.24)	0.21 (0.07-0.57)	0.30 (0.16-0.55)	0.53 (0.24-1.20)	1.23 (0.81-3.33)
Missing		0.42 (0.31-0.58)	0.59 (0.22-1.58)	0.66 (0.35-1.24)	0.58 (0.22-1.54)	1.19 (0.33-4.21)

* Adjusted for age, sex, calendar period, and baseline CD4 cell count in the analysis of any cancer, Kaposi sarcoma, non-Hodgkin lymphoma, and Hodgkin lymphoma. Adjusted for age, calendar period and baseline CD4 cell count in the analysis of cervical cancer. No adjustment was made in the analysis of leukaemia. Age was treated as a time-updated variable in all models.

† To preserve proportional hazards, we stratified the model for any cancer with respect to CD4 cell count and the model for cervical cancer with respect to age.

Figure 1: Hazard ratios (solid lines) with 95% confidence intervals (shaded areas) for cancer incidence, comparing a grid of baseline CD4 cell counts to the reference value of 200 cells/ μL . The models for any cancer, Kaposi sarcoma, non-Hodgkin lymphoma and Hodgkin lymphoma are adjusted for age, sex, and calendar period. The model for cervical cancer is adjusted for age and calendar period. The model for leukaemia is not adjusted for other variables.

